

CITY OF LAKE FOREST PARK
**URBAN FOREST ECOSYSTEM SERVICES
AND VALUES REPORT**

DECEMBER 2023

Acknowledgements

DCG/Watershed Contributors

Kim Frappier, Project Manager & Author
Sam Payne, Data Analyst & Author
Nathan Burroughs, GIS Analyst
Lars Freeman-Wood, Lead Field Researcher
Sage Yuasa, Field Researcher
Evan Earhart, Field Researcher
Drew Foster, Field Researcher & Analyst
Kenny Booth, Principal




City of Lake Forest Park

Phillip Hill, City Administrator
Larry Goldman, City Council
Steve Bennett, Planning Director
Hannah Swanson, City Arborist
Richard Olmstead, Tree Board Chair
Doug Sprugel, Tree Board
Marty Byrne, Tree Board
Sandra LeVar, Tree Board
Mark Phillips, Tree Board

We wish to express our gratitude to the i-Tree project team for their invaluable contribution in developing methodologies and software which facilitated the analysis in this study. Contributors to the i-Tree project are listed as the U.S. Forest Service, Davey Tree Expert Company, The Arbor Day Foundation, Society of Municipal Arborists, International Society of Arboriculture, and Casey Trees.

Project funding is from the City of Lake Forest Park.

Uncredited photos by Sam Payne, Katy Crandall, and Lexi Ochoa.





Summary

This report presents an evaluation of Lake Forest Park's urban and community forest through an i-Tree Eco plot sample inventory. Utilizing plot data obtained in 2022 and 2023, the i-Tree Eco model provides an assessment of urban forest health, structure, and threats as well as the ecosystem services and values trees provide the community. Additionally, tree canopy height is modeled using LiDAR to better understand the distribution of various canopy heights in the City's tree population. The following list summarizes key findings from this research effort.

- There are a total of 297,056 ($\pm 39,070$ SE) trees estimated to be in Lake Forest Park with a mean density of 129 trees per acre (TPA).
- Canopy cover is estimated at 50.6%.
- The most common tree species are Douglas-fir (16%), bigleaf maple (11%), western red cedar (9%), cherry laurel (8%), bitter cherry (6%), and English holly (6%). Of all trees, 63% are native to Washington.
- Less than 1% of trees are designated as noxious weeds in King County, however, 19% are listed as weeds of concern. The most abundant weeds of concern are cherry laurel, English holly, and sweet cherry.
- The age classification of trees trends youthful, with an abundance of smaller trees that will eventually replace the aging canopy. Trees less than 6-inches DBH account for 48.6% of the tree population.
- Leaf area density in the Large Residential stratum (parcels $> \frac{1}{4}$ acre) is three times greater than the Small Residential stratum (parcels $\leq \frac{1}{4}$ acre), and seven times greater than the Town Center stratum.
- The Lake Forest Park urban forest provides benefits valued at \$4.1 million annually for removing pollution, reducing runoff, sequestering carbon, and lowering energy usage.
- Carbon storage of the total urban forest is valued at \$16.6 million, and the replacement value is estimated at \$531 million.
- Of the 53 pests and pathogens that i-Tree assessed, 15 are present in King County. The economic impacts of these species are evaluated for each tree species and pest species.
- The canopy height model indicates that the proportion of tall trees, those greater than 135 feet in height, have increased by 21% from 2016 to 2021. The proportion of the tallest trees, those greater than 165 feet increased by 86% during this period, albeit accounting for less than 1% of the total tree population.



Table of Contents

INTRODUCTION	1
PROJECT BACKGROUND AND OBJECTIVES	1
SUMMARY OF URBAN FOREST BENEFITS	3
METHODS	4
I-TREE STUDY DESIGN	4
LIMITATIONS AND ASSUMPTIONS	5
CANOPY HEIGHT MODEL	6
RESULTS	7
TREE CHARACTERISTICS OF THE URBAN FOREST	7
URBAN FOREST COVER AND LEAF AREA	10
AIR POLLUTION	12
CARBON SEQUESTRATION AND STORAGE	13
SURFACE WATER RUNOFF	14
TREE BENEFITS SUMMARY	15
PESTS AND PATHOGENS	16
CANOPY HEIGHT MODEL RESULTS	19
DISCUSSION	20
CLIMATE ADAPTATION AND RESILIENCE	20
PROTECTION OF SIGNIFICANT AND LARGE TREES	21
INVASIVE SPECIES MANAGEMENT	22
ADDITIONAL CONSIDERATIONS	23
REFERENCES	24
APPENDICES	
APPENDIX I. I-TREE ECO MODEL AND FIELD MEASUREMENTS	
APPENDIX II. RELATIVE TREE EFFECTS	
APPENDIX III. COMPARISON OF URBAN FORESTS	
APPENDIX IV. GENERAL RECOMMENDATIONS FOR AIR QUALITY IMPROVEMENT	
APPENDIX V. TREE POPULATION	
APPENDIX VI. POTENTIAL RISK OF PESTS	



Photo provided by City of Lake Forest Park

Introduction

Lake Forest Park's urban and community forest consists of street trees, forested parks and open spaces, as well as trees on private residential, commercial, and industrial properties. These urban forest resources provide numerous ecosystem services, public health, and economic benefits to the people who live, work, and recreate here. Jurisdictions across King County and the State of Washington are faced with the need to support smart growth and development, environmental sustainability, and climate change resilience. Protecting green infrastructure such as tree canopies is critical to addressing these public and environmental health issues while ensuring the livability of Lake Forest Park. The first critical step to stewarding and managing this natural resource is understanding what we have.

The City of Lake Forest Park program has invested in tree inventories, canopy cover modeling, and studies investigating urban forest structure and values to guide the urban forestry program. This data has been used to inform and guide management actions, policies, municipal code updates, budget development, and identify

additional analysis needs. To date, the City has developed the following urban forest analysis and management plans:

- 2005 and 2016 Canopy Analyses
(LiDAR based studies)
- 2011 Urban Forest Effects and Values
(i-Tree Eco Analysis)
- 2010 Community Forest Management Plan

Project Background and Objectives


To build from the previous i-Tree Eco study published in 2011, the City contracted with DCG/Watershed in 2022 to conduct a follow-up survey to assess Lake Forest Park's community forest 10 years later. This analysis was first conducted in 2011 by the City arborist and the Lake Forest Park Tree Board, at that time a subset of the Environmental Quality Commission, with community volunteers participating in plot data collection.

The primary objectives of this 2022-2023 i-Tree study are to characterize urban forest structure and composition by collecting data on tree size, species, and health conditions. This data, along with other site level information within the specific study areas is then used to calculate the environmental and economic benefits at a city-scale.

Studying the structure and composition of the urban forest through the i-Tree analysis provides us with a more detailed understanding of Lake Forest Park's city-wide tree canopy, which were conducted in 2005 and again in 2016 using LiDAR analysis. Urban forest structure refers to the horizontal and vertical arrangement of trees, shrubs, and other plants, and their underlying abiotic environments, and is relevant to management because the physical arrangement in three-dimensional space influences the functions and ecosystem services provided by a forest. Composition refers to trees or other plant species that make up a forest.

Lake Forest Park's urban tree canopy covered 43% of the total City area in 2004, which became a baseline for forest management goals established in the City's 2010 Community Forest Management Plan. To reflect the diverse landscapes and development regulations within Lake Forest Park, canopy cover goals were established by land use types to be 50% in suburban residential area (lots $>1/4$ acres), 25% in urban residential area (lots $<1/4$ acres), and 15% in business districts. These were informed in-part by benchmarks recommended by American Forests. By 2016, total urban forest canopy increased to 50% based on a study by Elm (2016). This is comparable with recent analysis from i-Tree Landscape using high resolution data from 2017 which resulted in a canopy cover of 48%.

Canopy cover goals have been refined in Lake Forest Park Municipal Code Section 16.14.070 to represent



2016 Tree Canopy Coverage per Parcel
City of Lake Forest Park Tree Canopy Study

elm
Environmental
Management
Landscape
Urban Design
Urban Forest Studies

Figure 1. Lake Forest Park Tree Canopy Cover in 2016, reproduced from Elm (2016).

Table 1. Canopy coverage goals by zoning designation.

Zoning Designations	Canopy Coverage Goal
Single-family lots greater than 15,000 square feet	58%
Single-family lots 10,000 – 15,000 square feet	39%
Single-family lots less than 10,000 square feet	28%
Multifamily	15%
Commercial	15%
Southern Gateway Single Family	15%
Southern Gateway Transition	10%
Southern Gateway Corridor	5%

city zoning designations. As shown in Table 1, canopy cover goals range from 58% in zones with the lowest development intensity to 5% in zones with the greatest development intensity.

Summary of Urban Forest Benefits



Pollution Abatement: Urban forests serve as natural filters which improve water quality and air quality by trapping, absorbing, and transforming pollutants and excess nutrients, resulting in public health benefits, lower illness rates, and safeguarding ecosystems.



Shade and Cooling: Cities and metropolitan areas experience greater temperatures due to land use changes which alter the energy budget in an urban setting, known as the urban heat island effect. Through shading and evapotranspiration, urban forests mitigate the heat island effect through shading and cooling which lowers air and surface temperatures in densely populated regions.



Stormwater Reduction: Rainfall on impermeable surfaces, like concrete and asphalt, generates stormwater issues in cities, leading to problems such as flooding, water quality impairments, and reduced continuity of streamflow. In natural systems, rainwater interception and evapotranspiration minimize stormwater and reduce the reliance on costly engineered stormwater solutions.



Wildlife Habitat: Urban forests function as crucial wildlife habitats within the urban landscape, supporting a diverse range of species that have adapted to living alongside humans. These flora and fauna communities rely on these forests for essential resources, including refuge, food, water, and shelter, in an otherwise demanding environment.



Carbon Sequestration and Storage: Carbon dioxide (CO₂), the primary greenhouse gas driving global warming, is absorbed, and stored by trees during photosynthesis. This sequestered carbon is stored in the plant tissues during the lifetime of a tree.



Noise Buffering: Urban forests and tree canopies serve as natural noise buffers, reducing sound from traffic and other sources. The reduction of nuisance noise is beneficial to human health and well-being and can minimize noise impacts which negatively affect wildlife habitat.



Economic Benefits: Trees bring numerous economic advantages, such as higher property values, increased business traffic, heightened demand, tourism attraction, reduced energy costs, and resident appeal. Research indicates that urban forest programs typically yield substantial returns on investment, believed to be 2:1 or more (Endreny 2018).



Human Health and Wellness: Urban trees provide intangible yet significant societal benefits including recreation, enhancing the aesthetics of city streets, and fostering community pride and identity. Research also shows that trees play a role in improving health outcomes, reducing stress, enhancing mental well-being including cognition, attention, and anxiety, clinical outcomes, and crime reduction (Wolf et al. 2020).

Methods

i-Tree Study Design

The i-Tree Eco study was conducted using pre-stratified protocols to obtain representative samples with randomized 0.1-acre and 0.05-acre plots. Strata are consistent with the 2011 Lake Forest Park i-Tree study design for continuity in management units; these include parcels $>\frac{1}{4}$ -acres (Large Residential), parcels $\leq\frac{1}{4}$ -acres (Small Residential), and the commercial town center. Road networks are excluded from sample selection since they are interwoven amongst other strata and incorporated into calculations for total strata area.

Plots are located on both public and private lands. To secure permission to collect data on private parcels, the City arborist, with support from DCG/Watershed staff, contacted landowners via mail, email, the City newsletter, and door-knocking. Additional randomly selected plots were generated in instances where permission was not granted, until the required number of research plots was reached.

Data from 100 plots were collected in 2022 and 2023. An additional 60 plots were planned in the study design but could not be collected due to being denied access onto private property.

Once processed with the user defined data and configuration, i-Tree provides statistical analysis and actionable insights on a range of urban forestry topics including structure and composition, benefits and costs, air quality interactions, and pest analysis. Analysis of invasive species was conducted using information from the King County Noxious Weed Board, and species designations recorded in the i-Tree Eco software were disregarded.

i-Tree is a software suite and a set of tools developed by the USDA Forest Service and various partners to quantify the benefits and values of urban trees and forests. It provides a platform for assessing and managing urban forest ecosystems, focusing on the many environmental, economic, and societal benefits they offer.

i-Tree Manuals and Software Versions

- ❖ i-Tree Software Suite v6.0
- ❖ i-Tree Eco v6.0 User Manual
- ❖ i-Tree Eco v6.0 Field Manual

Stratum	Acres	Number of Plots
Town Center	19	8
$\leq\frac{1}{4}$ Acres	532	52
$>\frac{1}{4}$ Acres	1750	40





Figure 2. Strata and plot location map.



Urban Forest Measurements

This project utilizes data collection techniques as described in the i-Tree Eco v6.0 Field Manual. DCG/Watershed field researchers performed a range of measurements for each plot, encompassing both general plot characteristics and tree-specific measurement details. Plot-level and tree-level parameters are outlined in the graphic to the left. A total of 631 trees were assessed; this study includes all trees with a diameter at breast height (DBH) greater than one inch.



Plot Metrics


Plot ID	
Date	
Field Crew	
Plot Center Address	Distance to Plot Center
Coordinates (Lat/Long)	Direction from Plot Center
Tree Cover (%)	Tree Species
Shrub Cover (%)	DBH
Plantable Space (%)	Crown Condition (% Dieback)
Land Use	Tree Height
Ground Cover	Crown Top and Base Height
Comments	Crown Width (Bidirectional)

Tree Metrics

Tree ID	
Date	
Status	
Distance to Plot Center	
Direction from Plot Center	
Tree Species	
DBH	
Crown Condition (% Dieback)	
Tree Height	
Crown Top and Base Height	
Crown Width (Bidirectional)	
Percent of Crown Missing	
Crown Light Exposure	
Nearby Building Distance and Direction	
Street Tree	
Comments	

Limitations and Assumptions

Reported data were generated using i-Tree Eco, and therefore, limited by the associated model assumptions. Data provided by i-Tree Eco do not output standard error or other quantifiable metrics of sampling uncertainty for derived metrics. Standard error is reported for certain plot-level metrics supported by i-Tree Eco. Studies of i-Tree sampling methodology suggest that a 100-plot sample has an expected relative standard error (SE) of approximately 17%, however, this will differ by study and among assessed metrics (Nowak et al. 2008). Caution is advised in ascertaining trends between this study and the prior 2011 Lake Forest Park i-Tree Eco study for metrics which lack standard error metrics.




Light Detection and Ranging (LiDAR) can be used to provide highly accurate and spatially explicit models of urban forests. Canopy height models (CHM) are useful as a tool in urban forest management to quantify forest structure. Pictured (left) is a graphic depicting a CHM model of tree canopy height. Other LiDAR applications in forestry include canopy cover analysis, forest health assessment, biomass and carbon estimation, tree inventory mapping, and urban planning and design.

Canopy Height Model

This assessment includes a canopy height model (CHM) to provide information on urban forest structure and insight into retention of the City's largest and tallest trees. The CHM utilized LiDAR data from the two most recent LiDAR flights on publicly available databases, 2016 and 2021¹.

Modeling was completed in the R Program using the 'lidR' package, an open-source software integrated into the R ecosystem, for the purpose of manipulating and visualizing LiDAR data with applications in forestry. Canopy height model and tree top identification algorithms were used to identify tree heights with a variable search window. Trees overlapping buildings were removed from the model output using land classification data from the Washington Department of Natural Resources Urban Forestry's 2022 King County Land Cover Metrics dataset and outliers below 15 feet in height were removed because they could not reliably be distinguished from other shrubs or infrastructure.

This process yields a tree population point layer with canopy height attribute values that were evaluated for trends in canopy height over time.



¹ LiDAR Data obtained from the Washington Department of Natural Resources LiDAR Portal. Sourced information includes 2016 data from Quantum Spatial and 2021 data from the Washington Geologic Survey.

Results

Tree Characteristics of the Urban Forest

Lake Forest Park is estimated to contain 297,056 (\pm 39,070 SE) trees, which is greater than the reported population in 2011, with 249,000 trees. Canopy cover is estimated as 50.6% of the City area, ranking among the most heavily forested municipalities in the region. This is similar to other recent Lake Forest Park canopy cover estimates including the study conducted by Elm in 2016, which estimated canopy cover of 50%, and i-Tree Landscape, which estimated canopy cover of 48% in 2017 (data obtained from i-Tree Landscape in November 2023). While tree population and canopy cover appear to be increasing, the study methods do not support tests of statistical significance, and the error inherent in comparisons of this type do not permit us to say that there is statistical support for those trends.

Canopy cover is greatest in the Large Residential stratum (57%), followed by the Small Residential stratum (31%), and then the Town Center (13%). The defined strata in the study design do not allow for a direct comparison with the City's canopy cover goals separated by zone, though a side-by-side comparison in Table 2 displays those most closely related. Based on these estimates, the Large Residential and Small Residential strata meet the City's canopy cover goals while the Town Center lags slightly.




Figure 3. Canopy cover by stratum.




Figure 4. Tree density by stratum.

The average tree density in Lake Forest Park is estimated to be 129 trees per acre (TPA). Large Residential areas have the highest tree density, followed by Small Residential areas, and Town Center as shown in Figure 4.

Table 2. Comparison of canopy cover across i-Tree Eco study strata with city goals. Canopy cover goals for zoning designations do not correspond directly to the strata within the study design, and the nearest type is included for reference.

Stratum	City Goals	Corresponding Zones	Estimated Canopy Cover
Large Residential (>1/4 ac)	50%	Single-family lots 10,000 – 15,000 square feet Single-family lots >15,000 square feet	57%
Small Residential (<1/4 ac)	25%	Single-family lots <10,000 square feet	31%
Town Center	15%	Commercial	13%

There are 72 tree species represented in the study, though many other rare or infrequent species are likely present throughout Lake Forest Park which were not captured in the sampling plots. Since several tree species are not reliably identifiable to species in rapid field assessments, several of these taxa were identified to the genus level including certain apples (*Malus* spp.), plums (*Prunus* spp.), golden chain trees (*Laburnum* spp.), crape myrtles (*Lagerstroemia* spp.), privets (*Ligustrum* spp.), yews (*Taxus* spp.), and willows (*Salix* spp.). Since each of these likely represent multiple species, or hybrids of multiple species, overall species richness is assumed to be slightly higher than reported.

Species richness at the city-scale is greatest in the Small Residential stratum despite having a smaller total area than the Large Residential Stratum, with 54 and 42 species respectively (Figure 5).

Conversely, species richness at the local scale is greatest in the Large Residential stratum which has more species per acre than the Small Residential Stratum; with 22 and 11 species respectively. Both metrics of species richness are lowest in the Town Center stratum.

Douglas-fir, bigleaf maple, and western red cedar continue

to be the most common trees and are native to the Puget Lowlands Ecoregion. Diversity is key to resiliency in urban forests, particularly regarding impacts from disease and insects, and climate change.




Figure 5. Species richness and species per acres by strata.




Figure 6. Composition chart of the most abundant tree species.

Lake Forest Park has a greater canopy cover and tree density than any of the cities which i-Tree listed as comparable. Of these, Atlanta is reported to have the greatest tree canopy cover at 36.7% and Morgantown is reported to have the greatest tree density at 119 TPA.



Tree size class distributions provide a snapshot of forest structure that informs management strategies. Among these, it is useful to know whether a forest has a young or aging population. Currently, 71% of trees in Lake Forest Park are less than 12" DBH, indicating a skew toward younger or smaller trees.

Despite a youthful population, or Type 1 distribution (Morgenroth et al. 2020), the tree size class distribution skews slightly larger than the prior 2011 i-Tree Eco study. The percentage of the largest trees, those above 30" DBH, have increased since 2011 and now account for 5% of trees. Trees greater than 24" inches DBH now account for 10% of the total tree population, an increase from 2011.

Trees in Lake Forest Park are estimated to be 63% native to Washington overall, concentrated most highly in the Large Residential stratum, followed by the Small Residential and Town Center stratum (Figure 7).




Figure 8. Tree DBH class distribution by stratum.




Figure 7. Native status of trees by stratum.

Trees designated by King County or Washington State as noxious weeds comprise less than 1% of the tree population. These are represented by only one species, common hawthorn. However, 19% of trees are species listed by King County as weeds of concern. These include cherry laurel, sweet cherry, European mountain ash, black locust, horse chestnut, and English holly.




Figure 9. Tree DBH distribution.

Urban Forest Cover and Leaf Area

Leaf area density is greatest in the Large Residential stratum compared to other strata due to high tree density and the presence of larger trees. As a result, leaf area density in the Large Residential stratum is three times greater than the Small Residential stratum, and seven times greater than the Town Center stratum (Figure 8). A handful of species contribute most of the leaf area including Douglas-fir, western red cedar, and bigleaf maple (Table 3).

The importance value of each species represents the sum of the percent cover of a specific species and the leaf area percentage. This indicates which species dominate the urban canopy structure but are not always the best species to plant. The leaf area is an informative metric because it directly correlates with many urban forest functions and benefits such as avoided stormwater runoff.

Table 3. Leaf area, importance value, and percent of population by tree

Tree Species	Population (%)	Leaf Area (%)	Importance Value
Douglas-fir (<i>Pseudotsuga menziesii</i>)	16.3	35.5	51.8
Bigleaf maple (<i>Acer macrophyllum</i>)	11.1	18.9	30.0
Western red cedar (<i>Thuja plicata</i>)	9.1	15.5	24.6
Cherry laurel (<i>Prunus laurocerasus</i>)*	8.2	0.7	8.9
Red alder (<i>Alnus rubra</i>)	2.2	5.8	8.0
English holly (<i>Ilex aquifolium</i>)*	6.2	1.3	7.5
Vine maple (<i>Acer circinatum</i>)	5.2	1.9	7.1
Bitter cherry (<i>Prunus emarginata</i>)	6.3	0.7	6.9
Sweet cherry (<i>Prunus avium</i>)*	4.9	0.1	5.0
Deodar cedar (<i>Cedrus deodara</i>)	0.3	3.9	4.3
Western hemlock (<i>Tsuga heterophylla</i>)	2.8	1.3	4.1
Hinoki cypress (<i>Chamaecyparis obtusa</i>)	3.1	0.5	3.7
Japanese maple (<i>Acer palmatum</i>)	2.8	0.5	3.3
Western white pine (<i>Pinus monticola</i>)	0.4	2.8	3.2
Giant Sequoia (<i>Sequoiadendron giganteum</i>)	0.3	2.7	3.0
Arborvitae (<i>Thuja occidentalis</i>)	2.3	0.1	2.3
Black poplar (<i>Populus nigra</i>)	1.2	1.0	2.2
Portuguese laurel (<i>Prunus lusitanica</i>)	1.9	0.3	2.2
Sitka spruce (<i>Picea sitchensis</i>)	0.3	1.6	2.0
Pacific dogwood (<i>Cornus nuttallii</i>)	1.6	0.3	1.8
Plum (<i>Prunus</i> spp.)	1.3	0.1	1.4
Shore pine (<i>Pinus contorta</i>)	0.7	0.4	1.1
Blue spruce (<i>Picea pungens</i>)	0.4	0.7	1.1

* Designated as weed of concern by King County.



Total leaf area is defined as the one-sided area of all leaves in the study area. This differs from canopy cover because individual leaves may overlap within and among trees.



The total plantable space in Lake Forest Park is estimated to be 22.4% (± 2.9 SE), which represents opportunities for additional tree planting. This is defined as the amount of land area with suitable soils that are not under existing tree canopies or other overhead or land use restrictions that would prohibit tree planting (e.g., developed park or playfield).

Groundcover composition is a predictor variable for certain tree benefits estimated by i-Tree Eco because of the interaction between these benefits and ground-level processes. Stormwater avoidance, for example, is informed by the amount of impervious surface.

Groundcover composition is consistent with expectations for the land use types, with high intensity land uses having the most buildings and impervious surfaces, and the low intensity land uses having the most groundcover vegetation, duff/mulch, and bare soil. Impervious surfaces² are highest in the Town Center (82%), followed by Small Residential areas (51%), then Large Residential areas (25%).

The low building cover estimated in the Town Center is believed to be a result of a low sample size, since a significant portion of the site appears to be composed of buildings based on visual estimates. Where shrubs are present in sample plots, groundcover type below the shrub layer is recorded.




Figure 10. Leaf area density by stratum.




Figure 11. Ground cover composition by stratum.

² Impervious surfaces includes cement, tar/asphalt, other impervious surface, rock, and building.



Air Pollution

Many urban areas have high levels of air pollution which negatively impacts the health of humans and ecosystems. Urban forests mitigate the effects of air pollution through several processes including the absorption and particulate matter filtration, air temperature cooling, and reducing the energy consumption of buildings. While trees also emit volatile organic compounds (VOCs) that contribute to the formation of ozone (O_3), studies show that high tree cover is correlated with a reduction in ozone formation (Nowak and Dwyer 2000).

The Lake Forest Park urban forest canopy is estimated to remove 1,607 pounds of carbon monoxide (CO), 33,013 pounds of nitrogen dioxide (NO_2), 69,299 pounds of O_3 , 93,657 pounds of particulate matter less than 10 microns and greater than 2.5 microns (PM10), 12,458 pounds of particulate matter less than 2.5 microns (PM2.5), and 2,704 pounds of sulfur dioxide (SO_2) annually. This removal has an associated value of \$2.55 million.

Air pollution removal varies temporally, as shown in Figure 10. Some pollutants such as NO_2 and O_3 are removed at greater levels during the summer growing season while PM2.5 and PM10 removal is greatest during the fall and winter. Since some types of air pollution removal correlates with leaf area, the distribution of evergreen and deciduous trees also influences the magnitude of temporal variation. The large spike in PM10 in September and October is due to high concentrations present from wildfire smoke.




Figure 12. Estimated monthly pollution removal by the Lake Forest Park urban forest.




Figure 13. Estimated monetary value of air pollution removal annually.



Carbon Sequestration and Storage

Tree canopy cover in Lake Forest Park is not just a local issue. Global climate change is largely driven by carbon dioxide (CO₂) emissions, a compound which trees uptake and sequester during photosynthesis. Carbon is stored in tree leaves and woody tissues, and therefore, reduces the amount of atmospheric carbon otherwise contributing to climate change. Carbon will remain in a tree until it eventually decomposes, where it may either be released to the atmosphere, returned to soil, or absorbed by other organisms.

The Lake Forest Park urban forest is estimated to remove 2,639 tons of carbon annually. Areas with greatest tree cover also provide the greatest levels of CO₂ sequestration, such as the Large Residential

stratum, which provide two to six-fold more than the other strata on a per-area basis (Figure 14). The estimated value of this benefit is \$450,000 per year. Carbon storage and carbon sequestration values are based on the societal costs of carbon emission such as climate change impacts on flooding, sea level rise, and agriculture (Appendix I).

Carbon storage is also valuable to quantify because a tree that decomposes will eventually release CO₂ back into the atmosphere. Trees in Lake Forest Park collectively store 97,300 tons of carbon, with an estimated value of \$16.6 million. Douglas-fir, bigleaf maple, western red cedar, and black poplar are the tree species which currently have the greatest amount of carbon storage.




Figure 14. Carbon sequestration each year by stratum.




Figure 15. Total carbon storage by stratum.

Climate change is the process of shifting global and regional climate patterns, driven primarily by anthropogenic activities such as fossil fuel emissions and deforestation. These result in increased concentrations of greenhouse gases in the atmosphere which lead to globally rising temperatures, altered weather patterns, and sea level rise, which affect societies, economies, and ecosystems across the planet. Changing climates also mean cities need to manage for resilient forests which can tolerate shifting conditions.



Surface Water Runoff

Runoff from impermeable surfaces is a significant source of water pollution and flooding, posing risks to both human and environmental well-being while imposing substantial economic costs. Trees play a role in mitigating runoff through evapotranspiration, a combination of processes which include the interception of rainwater, evaporation, and transpiration, and thereby, return water to the atmosphere. Additionally, trees enhance the ability of rainwater to infiltrate into soils through inputs of organic matter and improving porosity. The combination of these processes results in the attenuation of pollution laden runoff and reduction in the severity of flooding events. Urban forests in Lake Forest Park are estimated to reduce runoff by 50.4 million gallons per year.

Urban forests also reduce the need for cities to rely on costly built infrastructure to manage water quality and quantity issues. This “green infrastructure” is estimated to provide Lake Forest Park with estimated economic benefit at \$450,000 per year for water quality and flood reduction benefits they provide. The majority of these benefits are provided in the Large Residential stratum, where tree density and leaf area are greatest.




Figure 16. Avoided runoff per annum, by stratum.




Figure 17. Value of avoided runoff per annum, by stratum.



Tree Benefits Summary

The total economic benefit of trees in Lake Forest Park is estimated to be \$4.1 million per year, when accounting for energy savings, gross carbon sequestration, pollution removal, and avoided runoff (Table 4). On an individual basis, this amounts to \$13.79 per tree on average.

Tree replacement values are another useful measure when managing forests since it is more expensive to replace trees than preserve existing trees. The collective replacement value of all trees in Lake Forest Park is estimated to be \$531 million in addition to the \$16.9 million provided by carbon storage. The high cost of tree removal can inform public policy and management decisions regarding tree preservation and replacement on public and private land.

Trees in Lake Forest Park also generate 5,316 tons of oxygen every year, however, this benefit is believed to be relatively insignificant due to the vast reserves of oxygen in the atmosphere and production from oceanic systems (Broecker 1970; i-Tree 2023).

Table 4. Total Lake Forest Park tree benefits summary.

Benefits	Annual Value	Annual Value Per Tree
Energy & Carbon Emission Reduction	\$646,683	\$2.17
Gross Carbon Sequestration	\$450,102	\$1.53
Pollution Removal	\$2,545,703	\$8.57
Avoided Runoff	\$450,254	\$1.52
Total Benefits	\$4,092,742	\$13.79

Urban forests result in a net reduction in energy use through shading, evaporative cooling, and blocking of winter winds which are estimated to save Lake Forest Park residents \$542,683 per year. Additionally, the value of reduced carbon emissions resulting from energy savings is valued at \$104,000 per year.



Pests and Pathogens

Trees are susceptible to pests and pathogens that are capable of impacting tree viability, resulting in reduced lifespan, hazard conditions and sometimes mortality. The i-Tree Eco model includes an analysis of the susceptibility of Lake Forest Park's urban forests to 53 common pests and pathogens to evaluate risks and management priorities. While some pests and pathogens are naturally occurring and play an important role in forest ecological processes, others may have significant negative ecological and economic impacts. This section introduces the types of pests and pathogens identified through the i-Tree analysis based on data from the Forest Health Technology Enterprise Team (i-Tree 2023). See Appendix V for the complete list of pests and pathogens assessed through i-Tree. As described in Figure 18, the three species that could impact the highest percentage of canopy species in Lake Forest Park are two fungal pathogens and one




Figure 18. Susceptibility by trees to the 15 evaluated pests and pathogens which are currently known to be present in King County.

insect commonly found in Pacific Northwest forests (Armillaria root disease, Heterobasidion root disease, and western spruce budworm).

Armillaria Root Disease (*Armillaria* sp.) refers to a group of fungi that causes reduced leader growth and foliage discoloration and thinning, spreading through a tree's root system (Allen et al. 1996).

Pests and pathogens included in the i-Tree analysis have been documented within King County limits but does not confirm their presence in the trees surveyed within the study plots. This research did not include an advanced level of tree health analysis beyond the standard i-Tree data collection protocols.

Heterobasidion Root Disease (HRD; *Heterobasidion annosum*, *H. occidentale*), also called Annosus root and butt rot, is a fungus known to impact native conifers as well as bigleaf maple and red alder. In younger trees, symptoms include a reduction in the leader and branch growth, chlorotic foliage and a distressed cone crop. (Allen et al. 1996).



Photo by USDA Forest Service; fruiting body of *H. occidentale*.

Western spruce budworm (*Choristoneura occidentalis*, *C. freemani*) is an insect native to western North America and is a widespread defoliator of several native conifer species. It feeds upon and defoliates Douglas-firs, spruce, and the true firs (e.g., white fir and subalpine fir). The larvae feed on the current year's needles and buds giving the canopy a red-brown or grayish appearance with thinning foliage and produces a new generation annually.



Photo by: Montana State University Extension; Western spruce budworm.

Emerging Threats in Western Washington

Disease and pest outbreaks have increased in number and frequency in recent years due to international trade, travel, and climatic changes. New pests are introduced outside of their native range into ecosystems that have not evolved with the pest to develop any resistance. The effects of climate change, such as increases in seasonal and average air temperatures, increases in extreme heat, and prolonged drought, add abiotic stressors, weakening a tree's ability to defend against these diseases and pest pressures (Mauger et al. 2015). Additional pests and pathogens that could have an impact on canopy trees within Lake Forest Park include sooty bark disease, bronze birch borer, emerald ash borer, and non-native long-horned beetle species.

Sooty bark disease (*Cryptostroma corticale*) causes dieback primarily in maple species. To date, the fungus has been found to cause damage in sycamore maples (*Acer pseudoplatanus*), red maple (*A. rubrum*), Japanese maple (*A. palmatum*), vine maple (*A. circinatum*), and bigleaf maple (*A. macrophyllum*) in the Puget Sound region. The fungus infects the tree's vascular system and thrives during hot summers, proliferating in drought-stressed trees (Brooks et al. 2022).


Bronze birch borer (*Agrilus anxius*) is a beetle whose larvae tunnel into live wood, creating extensive galleries leading to branch or trunk girdling, ultimately cutting the rest of the branch off from resources. Bronze birch borers are attracted to trees weakened by environmental stressors, age, or other diseases and pests (Antonelli 2008).


Emerald ash borer (*Agrilus planipennis*) has been present in the United States since 2002 but only recently has been confirmed in the Pacific Northwest Region as of 2022, where it was discovered in Oregon. While it has not yet been

sighted in the Puget Sound region, its spread into Washington State is expected. The emerald ash borer infects native and non-native ash trees (*Fraxinus* spp.). Like other borers, its larvae create extensive galleries, causing limb and trunk dieback leading to decline and eventual tree death (Bliss-Ketchum et al. 2021). Although ash trees were found to only compose a tiny fraction of the tree population (<0.1%), it is possible that more are present and were not represented in the samples. Oregon ash (*Fraxinus latifolia*) is a native tree which can be found near water or in wetland areas. Ashes are also commonly planted as street trees and as ornamentals in yards and gardens.



Asian, citrus, and red-necked long-horned beetles (*Anoplophora glabripennis*, *A. chinensis*, and *Aromia bungii* respectively) feed on the wood of hardwood trees. Although there are no known established populations of these beetles in Washington, they have reached local nurseries where they were eradicated. With continued global movement within the nursery trade, Washington will need to continually monitor these species. The beetles typically feed on both healthy and dying trees and are known to impact 40 host species including maples, horse chestnuts, willows, birches, and elms. There are locally known native look-a-likes which present a challenge to identification for nonprofessionals (WISC 2017). The Washington Invasive Species Council has resources for identifying the potentially invasive versus native beetles in King County.





Canopy Height Model Results

The distribution of tree heights in Lake Forest Park reveals that the proportion of tall trees, those greater than 135 feet in height, has increased by 21% in 2021 compared to 2016 (Figure 19). The proportion of the tallest trees, those greater than 165 feet has increased by 86% during this period, albeit accounting for less than 1% of the total tree population. This suggests that most tall trees are being retained, and that other small and moderate size trees are aging into the larger height classifications. The tallest tree is estimated to be 195 feet tall.

Trends of smaller trees vary by height class, although the proportion of trees in the moderate

height classes have tended to decrease while the smallest, those between 15-30 feet, are approximately equal. Since trees below 15 feet were removed in this analysis, plot samples collected as part of the i-Tree Eco inventory provide better insight into age distribution and forest regeneration. The canopy height model is less selective than the plot sampling method in finding smaller trees and subcanopy trees, so interpretations of age and regeneration are not as precise as other sampling methods. However, this analysis provides us with additional insight into the distribution of trees within the assessed range between 15 and 195 feet.




Figure 19. Histogram of tree heights in 2016 and 2021 based on the CHM.



Photo by Cori Whitaker

Discussion

The results of this i-Tree Eco study and canopy height model provide insight on the current composition and structure of Lake Forest Park's urban and community forest as well as quantify ecosystem service benefits and values. The results suggest a net increase in urban tree canopy cover and tree density during the last ten years and an increasing trend in the presence of large canopy trees, primarily comprised of Pacific Northwest native species.

The data obtained by this study provides City urban forest managers with practical information that is useful to develop urban forest management strategies and policies. Cities across the Puget Sound region and the Pacific Northwest face several challenges to steward resilient, regenerative, and viable urban forests. These include shifts in climate conditions, threats from current and emerging pests and pathogens, the potential for increases in urban wildfires, and continued development needed to meet regional housing needs. Urban forest managers are also tasked with ensuring that tree canopy remains equitably distributed throughout the City and that more densely developed land use zones have adequate green infrastructure to manage stormwater, minimize urban heat islands, provide shade, and foster both ecological health as well as human health and wellness.

Climate Adaptation and Resilience

Within the field of urban forest management, arborists, ecologists, foresters, and land managers continue to evaluate best management practices and adapt arboricultural strategies to the on-the-ground conditions impacting the resilience of urban forests.

Western Washington is expected to experience increasingly drier conditions and higher temperatures during the summer months, with potential increases in precipitation during the winter months (Mauger et al. 2015). This will present new and exacerbate current stressors on existing urban forests such as drought, insect and tree disease outbreaks, competition with invasive plant species, habitat loss and fragmentation, erosion, and wildfires. These stressors also create challenges for establishing the next generation of urban forest canopy, especially coupled with development pressures and the need to respond to the rising necessity for sustainable and affordable housing.

One strategy for establishing resilience within the urban forest is to increase tree diversity (at the family, genus, and species level) and ensure installed trees are climate adapted to current and future stressors, such as drought. Since most biotic and abiotic stressors exhibit variable effects among tree species, a diverse forest acts as an insurance



policy that minimizes risk from impacts to individual taxa.

The City of Lake Forest Park currently has an approved tree list which includes species that are appropriate for the built environment in LFP and that are drought tolerant. This is an important educational and management tool that should be periodically evaluated and updated to account for updated research and recommendations from the arboriculture and horticultural trades and account for climate resilience.

Protection of Significant and Large Trees

The tree size class distributions outlined in this study tell us that 71% of the City's forests are less than 12" diameter-at-breast-height (DBH), but that the percentage of large diameter trees (those greater than 24" DBH) has continued to increase during the last decade. Tree diameter correlates with tree height and volume and can be used as a metric to describe overall tree size and identify large trees, which are a management priority for the City. Large trees provide greater levels of ecosystem services such as stormwater capture and infiltration, cooling, and water quality improvements compared to small trees; and therefore, societal benefits are optimized when they are retained.

Since the majority of Lake Forest Park's urban forest is located on private residential, commercial, and industrial property, protection of significant and large diameter trees on privately owned property will be an important strategy as the City seeks to protect its existing tree canopy. The City currently regulates trees during development of private property through its tree ordinance – Chapter 16.14 *Tree Canopy Preservation and Enhancement* - as well as trees within shoreline jurisdiction through Chapter 16.18 *Shoreline Master Program*. These regulations prioritize the retention and protection of existing



trees and groves as well as replanting with new trees when removal is unavoidable due to tree risks, site development design, and storm damage.

The findings within this report can be used as a tool to educate and engage community members, private landowners, and the development community to encourage early assessment and integration of existing significant and exceptional trees in the pre-design or early design phase of new development. Another critical component is ensuring not only the long-term viability of a retained tree but ensuring that replacement trees are chosen using the “right plan, right place” approach, have adequate growing conditions (e.g., soil volumes, planter strip widths etc.) to reach maturity without impacting required infrastructure such as sidewalks, driveways, and utilities.

Invasive Species Management

Invasive plants are self-propagating and aggressive introduced species that are known to outcompete our native flora. They present significant challenges to urban forest health and management, including the economic investment for control and eradication as well as the costs to replace the ecosystem service benefits, they provided. This study provides an inventory of invasive trees, although shrub and understory components of the urban forest are not a part of the study design.

It is advantageous, both ecologically and economically, to control invasive plants before they

become widely established in urban forests. Once a forest is overrun by invasive species, there is often an associated drop in wildlife richness, as they do not support the same level of habitat structure or food availability as native ecosystems.

For example, two shrub species that are heavily impacting Lake Forest Park’s urban forest are Himalayan blackberry (*Rubus bifrons*) and English ivy (*Hedera helix*). Both species are Class C non-regulated weeds on the King County Noxious Weed List, are widely distributed throughout the Puget Sound region, and recommended for control where feasible due to their impacts (King County Noxious Weed Control Board 2023).

Himalayan blackberry is a woody shrub that forms dense brambles in clearings and forest understories that become so thick that no other plants will grow. Seed distribution is bird-facilitated which provides a constant source to new areas, so it is difficult to eliminate entirely. English ivy forms an expansive mat through the trunks and canopies of trees, and on the forest floor, that competes with native plants for nutrients and light. English ivy can significantly reduce a trees lifespan through this direct competition, by girdling the stem, and increasing limb and tree failure from the extra weight. Once invasive plants take hold, native tree seedling establishment is suppressed, halting the cycle of forest regeneration. Existing trees eventually senesce and fall, leaving gaps in the canopy that can be quickly filled in by the established invasive plants.



Of the trees surveyed, 19% are either listed as a Class C noxious weed (common hawthorn) or a Weed of Concern (sweet cherry, black locust, cherry laurel, English holly, and European mountain ash) by the King County Noxious Weed Program. Cherry laurel and English holly, black locust, and European mountain ash are widely used as ornamental landscape plants. However, these species compete with our native flora and naturalize in open spaces and critical areas. Stokes et al (2014) examined the dispersal, spread, and impact of English holly in St. Edwards State Park and found that native vegetation was greatly reduced under holly canopy. The study also reports that the holly population was spreading rapidly both through seed dispersal and vegetatively through expansion of tree clumps.

A potential strategy to address this problem could be for the City to develop a prohibited species list and other educational materials to discourage property owners and developers from introducing invasive shrub and tree species into new plantings. In addition, the City could consider implementing removal of both invasive shrub and tree species from public open spaces, replacing them with native tree and shrub species.

Additional Considerations

Scientific studies as well as programmatic and policy audits provide important data to evaluate the success of urban forest resource management strategies. In addition to the continuation of tree benefit analyses, the evaluation of other policies and regulations can inform municipal code updates, the effectiveness of current community education and outreach efforts, and additional support needed from community members in managing trees on their properties. Continued study of on-the-ground conditions coupled with evaluation of existing policies and best practices will provide the City with the tools and information needed to manage the valuable urban forest resource effectively and adaptively.

References

Allen, E.A., D.J. Morrison, and G.W. Wallis. (1996). *Common Tree Diseases of British Columbia*. Natural Resources Canada/Canadian Forest Service.

Antonelli, A. (2008). *Bronze Birch Borer* (Report No. EB1380E). Washington State University Extension. EB1380E. <https://pubs.extension.wsu.edu/bronze-birch-borer>

Brooks, B., Hulbert, J.M., Omdal, D., Elliot, M., and Chastagner, G. A. (2022). *Sooty bark disease diagnostic guide* (Report No. FS375E). Washington State University Extension. <https://pubs.extension.wsu.edu/sooty-bark-disease-diagnostic-guide>

Bliss-Ketchum L.B., R. Draheim, M. Hepner, and O. Guethling. (2021). *Emerald Ash Borer: Readiness and Response Plan for Oregon*. EAB Readiness and Response Steering Committee. <https://www.oregon.gov/odf/Documents/forestbenefits/eab-readiness-and-response-plan-for-oregon.pdf>.

Broecker, W. S. (1970). A boundary condition on the evolution of atmospheric oxygen. *Journal of Geophysical Research*, 75(18), 3553–3557. <https://doi.org/10.1029/JC075i018p03553>

Elm. (2016). *Tree Canopy Extent*, City of Lake Forest Park Tree Canopy Study.

Endreny, T. A. (2018). Strategically growing the urban forest will improve our world. *Nature Communications*, 9(1), 1160–1163. <https://doi.org/10.1038/s41467-018-03622-0>

City of Lake Forest Park. (2010). *Community Forest Management Plan*. <https://www.cityoflfp.gov/DocumentCenter/View/6175/Community-Forest-Management-Plan?bidId=>

City of Lake Forest Park. (2011). *I-Tree Ecosystem Analysis: Urban Forest Effects and Values*. i-Tree Report. <https://www.cityoflfp.gov/DocumentCenter/View/385/itreereport?bidId=>

City of Lake Forest Park. (n.d.). *General tree list for lake forest park, WA*. Accessed November 2023. <https://www.cityoflfp.gov/DocumentCenter/View/224/Tree-List?bidId=>

City of Lake Forest Park. (2023a). *Municipal Code Chapter 16.14 Tree Canopy Preservation and Enhancement*. Accessed October 2023. <https://www.codepublishing.com/WA/LakeForestPark/>

City of Lake Forest Park. (2023b). *Municipal Code Chapter 16.18 Shoreline Master Program*. Accessed October 2023. <https://www.codepublishing.com/WA/LakeForestPark/>

i-Tree. (2021a). *Field Guide*. <https://www.itreetools.org/documents/274/EcoV6.FieldManual.2021.10.06.pdf>

i-Tree. (2021b). *User's Manual*. https://www.itreetools.org/documents/275/EcoV6_UsersManual.2021.09.22.pdf

i-Tree. (2023). i-Tree Eco Software Suite v6.0.

King County Noxious Weed Control Board. (2023). *King County Noxious Weed List*. https://your.kingcounty.gov/dnrp/library/water-and-land/weeds/weedlists/2023-Official-Weed-List_common-name.pdf

Mauger, G.S., J.H. Casola, H.A. Morgan, R.L. Strauch, B. Jones, B. Curry, T.M. Busch Isaksen, L. Whitely Binder, M.B. Crosby, and A.K. Snover, 2015. *State of Knowledge: Climate Change in Puget Sound Report* prepared for the Puget Sound Partnership and the National Oceanic and Atmospheric Administration. Climate Impacts Group, University of Washington.

Morgenroth, J., Nowak, D. J., and Koeser, A. K. (2020). DBH Distributions in America's Urban Forests-An Overview of Structural Diversity. *Forests*, 11(2), 135-. <https://doi.org/10.3390/f11020135>

Nowak, D. J., and Dwyer, J. F. (2000). *Understanding the Benefits and Costs of Urban Forest Ecosystems*. In Handbook of Urban and Community Forestry in the Northeast (pp. 11–25). Springer US. https://doi.org/10.1007/978-1-4615-4191-2_2

Nowak, D., Walton, J., Stevens, J., Crane, D., & Hoehn, R. (2008). Effect of Plot and Sample Size on Timing and Precision of Urban Forest Assessments. *Arboriculture & Urban Forestry*, 34(6), 386–390. <https://doi.org/10.48044/jauf.2008.052>

Fellin, D.G., and Dewey, J.E. (1986). *Western Spruce Budworm* (Forest Insect and Diseases Leaflet 53). U.S. Forest Service. https://www.fs.usda.gov/Internet/FSE_DOCUMENTS/fsbdev2_043445.pdf

Stokes, D.L., Church, E.D., Conkright, D.M., and Lopiz, S. (2014). *Pictures of an Invasion: English Holly (Ilex aquifolium) in a Semi-Natural Pacific Northwest Forest*. Northwest Science, 88(2): 75–93.

Washington Invasive Species Council (WISC). (2017a). *Citrus, Asian, and Red-Necked Longhorned Beetles Fact Sheet*. <https://invasivespecies.wa.gov/wp-content/uploads/2019/07/InvasiveLonghornedBeetles-FactSheet.pdf>

Washington Invasive Species Council (WISC). (2017b). *Emerald Ash Borer, Stop the Invasion*. <https://invasivespecies.wa.gov/wp-content/uploads/2019/07/EmeraldAshBorer-FactSheet.pdf>

Wolf, K. L., Lam, S. T., McKeen, J. K., Richardson, G. R. A., van den Bosch, M., & Bardekjian, A. C. (2020). Urban Trees and Human Health: A Scoping Review. *International Journal of Environmental Research and Public Health*, 17(12), 4371-. <https://doi.org/10.3390/ijerph17124371>

Appendix I. i-Tree Eco Model and Field Measurements

i-Tree Eco is designed to use standardized field data from randomly located plots and local hourly air pollution and meteorological data to quantify urban forest structure and its numerous effects (Nowak and Crane 2000), including:

- Urban forest structure (e.g., species composition, tree health, leaf area, etc.).
- Amount of pollution removed hourly by the urban forest, and its associated percent air quality improvement throughout a year.
- Total carbon stored and net carbon annually sequestered by the urban forest.
- Effects of trees on building energy use and consequent effects on carbon dioxide emissions from power sources.
- Replacement value of the forest, as well as the value for air pollution removal and carbon storage and sequestration.
- Potential impact of infestations by pests, such as Asian longhorned beetle, emerald ash borer, gypsy moth, and Dutch elm disease.

Typically, all field data are collected during the leaf-on season to properly assess tree canopies. Typical data collection (actual data collection may vary depending upon the user) includes land use, ground and tree cover, individual tree attributes of species, stem diameter, height, crown width, crown canopy missing and dieback, and distance and direction to residential buildings (Nowak et al 2005; Nowak et al 2008).

During data collection, trees are identified to the most specific taxonomic classification possible. Trees that are not classified to the species level may be classified by genus (e.g., ash) or species groups (e.g., hardwood). In this report, tree species, genera, or species groups are collectively referred to as tree species.

Tree Characteristics:

Leaf area of trees was assessed using measurements of crown dimensions and percentage of crown canopy missing. In the event that these data variables were not collected, they are estimated by the model.

An analysis of invasive species is not available for studies outside of the United States. For the U.S., invasive species are identified using an invasive species list (Oregon Invasive Species Council 2014) for the state in which the urban forest is located. These lists are not exhaustive and they cover invasive species of varying degrees of invasiveness and distribution. In instances where a state did not have an invasive species list, a list was created based on the lists of the adjacent states. Tree species that are identified as invasive by the state invasive species list are cross-referenced with native range data. This helps eliminate species that are on the state invasive species list, but are native to the study area.

Air Pollution Removal:

Pollution removal is calculated for ozone, sulfur dioxide, nitrogen dioxide, carbon monoxide, particulate matter less than 2.5 microns, and particulate matter less than 10 microns and greater than 2.5 microns. PM2.5 is generally more relevant in discussions concerning air pollution effects on human health.

Air pollution removal estimates are derived from calculated hourly tree-canopy resistances for ozone, and sulfur and nitrogen dioxides based on a hybrid of big-leaf and multi-layer canopy deposition models (Baldocchi 1988; Baldocchi et al 1987). As the removal of carbon monoxide and particulate matter by vegetation is not directly related to transpiration, removal rates (deposition velocities) for these pollutants were based on average measured values from the literature (Bidwell and Fraser 1972; Lovett 1994) that were adjusted depending on leaf phenology and leaf area. Particulate removal incorporated a 50 percent resuspension rate of particles back to the atmosphere (Zinke 1967). Recent updates (2011) to air quality modeling are based on improved leaf area index simulations, weather and

pollution processing and interpolation, and updated pollutant monetary values (Hirabayashi et al 2011; Hirabayashi et al 2012; Hirabayashi 2011).

Trees remove PM2.5 and PM10* when particulate matter is deposited on leaf surfaces (Nowak et al 2013). This deposited PM2.5 and PM10* can be resuspended to the atmosphere or removed during rain events and dissolved or transferred to the soil. This combination of events can lead to positive or negative pollution removal and value depending on various atmospheric factors. Generally, PM2.5 and PM10* removal is positive with positive benefits. However, there are some cases when net removal is negative or resuspended particles lead to increased pollution concentrations and negative values. During some months (e.g., with no rain), trees resuspend more particles than they remove. Resuspension can also lead to increased overall PM2.5 and PM10* concentrations if the boundary layer conditions are lower during net resuspension periods than during net removal periods. Since the pollution removal value is based on the change in pollution concentration, it is possible to have situations when trees remove PM2.5 and PM10* but increase concentrations and thus have negative values during periods of positive overall removal. These events are not common, but can happen.

For reports in the United States, default air pollution removal value is calculated based on local incidence of adverse health effects and national median externality costs. The number of adverse health effects and associated economic value is calculated for ozone, sulfur dioxide, nitrogen dioxide, and particulate matter less than 2.5 microns using data from the U.S. Environmental Protection Agency's Environmental Benefits Mapping and Analysis Program (BenMAP) (Nowak et al 2014). The model uses a damage-function approach that is based on the local change in pollution concentration and population. National median externality costs were used to calculate the value of carbon monoxide removal (Murray et al 1994).

For international reports, user-defined local pollution values are used. For international reports that do not have local values, estimates are based on either European median externality values (van Essen et al 2011) or BenMAP regression equations (Nowak et al 2014) that incorporate user-defined population estimates. Values are then converted to local currency with user-defined exchange rates.

For this analysis, pollution removal value is calculated based on the prices of \$1,397 per ton (carbon monoxide), \$4,926 per ton (ozone), \$613 per ton (nitrogen dioxide), \$181 per ton (sulfur dioxide), \$330,079 per ton (particulate matter less than 2.5 microns), \$6,565 per ton (particulate matter less than 10 microns and greater than 2.5 microns).

Carbon Storage and Sequestration:

Carbon storage is the amount of carbon bound up in the above-ground and below-ground parts of woody vegetation. To calculate current carbon storage, biomass for each tree was calculated using equations from the literature and measured tree data. Open-grown, maintained trees tend to have less biomass than predicted by forest-derived biomass equations (Nowak 1994). To adjust for this difference, biomass results for open-grown urban trees were multiplied by 0.8. No adjustment was made for trees found in natural stand conditions. Tree dry-weight biomass was converted to stored carbon by multiplying by 0.5.

Carbon sequestration is the removal of carbon dioxide from the air by plants. To estimate the gross amount of carbon sequestered annually, average diameter growth from the appropriate genera and diameter class and tree condition was added to the existing tree diameter (year x) to estimate tree diameter and carbon storage in year x+1.

Carbon storage and carbon sequestration values are based on estimated or customized local carbon values. For international reports that do not have local values, estimates are based on the carbon value for the United States (U.S. Environmental Protection Agency 2015, Interagency Working Group on Social Cost of Carbon 2015) and converted to local currency with user-defined exchange rates.

For this analysis, carbon storage and carbon sequestration values are calculated based on \$171 per ton.

Oxygen Production:

The amount of oxygen produced is estimated from carbon sequestration based on atomic weights: net O₂ release (kg/yr) = net C sequestration (kg/yr) × 32/12. To estimate the net carbon sequestration rate, the amount of carbon sequestered as a result of tree growth is reduced by the amount lost resulting from tree mortality. Thus, net carbon sequestration and net annual oxygen production of the urban forest account for decomposition (Nowak et al 2007). For complete inventory projects, oxygen production is estimated from gross carbon sequestration and does not account for decomposition.

Avoided Runoff:

Annual avoided surface runoff is calculated based on rainfall interception by vegetation, specifically the difference between annual runoff with and without vegetation. Although tree leaves, branches, and bark may intercept precipitation and thus mitigate surface runoff, only the precipitation intercepted by leaves is accounted for in this analysis.

The value of avoided runoff is based on estimated or user-defined local values. For international reports that do not have local values, the national average value for the United States is utilized and converted to local currency with user-defined exchange rates. The U.S. value of avoided runoff is based on the U.S. Forest Service's Community Tree Guide Series (McPherson et al 1999; 2000; 2001; 2002; 2003; 2004; 2006a; 2006b; 2006c; 2007; 2010; Peper et al 2009; 2010; Vargas et al 2007a; 2007b; 2008).

For this analysis, avoided runoff value is calculated based on the price of \$0.01 per gallon.

Building Energy Use:

If appropriate field data were collected, seasonal effects of trees on residential building energy use were calculated based on procedures described in the literature (McPherson and Simpson 1999) using distance and direction of trees from residential structures, tree height and tree condition data. To calculate the monetary value of energy savings, local or custom prices per MWH or MBTU are utilized.

For this analysis, energy saving value is calculated based on the prices of \$96.70 per MWH and \$10.65 per MBTU.

Replacement Values:

Replacement value is the value of a tree based on the physical resource itself (e.g., the cost of having to replace a tree with a similar tree). Replacement values were based on valuation procedures of the Council of Tree and Landscape Appraisers, which uses tree species, diameter, condition, and location information (Nowak et al 2002a; 2002b). Replacement value may not be included for international projects if there is insufficient local data to complete the valuation procedures.

Potential Pest Impacts:

The complete potential pest risk analysis is not available for studies outside of the United States. The number of trees at risk to the pests analyzed is reported, though the list of pests is based on known insects and disease in the United States.

For the U.S., potential pest risk is based on pest range maps and the known pest host species that are likely to experience mortality. Pest range maps for 2012 from the Forest Health Technology Enterprise Team (FHTET) (Forest Health Technology Enterprise Team 2014) were used to determine the proximity of each pest to the county in which

the urban forest is located. For the county, it was established whether the insect/disease occurs within the county, is within 250 miles of the county edge, is between 250 and 750 miles away, or is greater than 750 miles away. FHTET did not have pest range maps for Dutch elm disease and chestnut blight. The range of these pests was based on known occurrence and the host range, respectively (Eastern Forest Environmental Threat Assessment Center; Worrall 2007).

Relative Tree Effects:

The relative value of tree benefits reported in Appendix II is calculated to show what carbon storage and sequestration, and air pollutant removal equate to in amounts of municipal carbon emissions, passenger automobile emissions, and house emissions.

Municipal carbon emissions are based on 2010 U.S. per capita carbon emissions (Carbon Dioxide Information Analysis Center 2010). Per capita emissions were multiplied by city population to estimate total city carbon emissions.

Light duty vehicle emission rates (g/mi) for CO, NOx, VOCs, PM10, SO2 for 2010 (Bureau of Transportation Statistics 2010; Heirigs et al 2004), PM2.5 for 2011-2015 (California Air Resources Board 2013), and CO2 for 2011 (U.S. Environmental Protection Agency 2010) were multiplied by average miles driven per vehicle in 2011 (Federal Highway Administration 2013) to determine average emissions per vehicle.

Household emissions are based on average electricity kWh usage, natural gas Btu usage, fuel oil Btu usage, kerosene Btu usage, LPG Btu usage, and wood Btu usage per household in 2009 (Energy Information Administration 2013; Energy Information Administration 2014)

- CO2, SO2, and NOx power plant emission per kWh are from Leonardo Academy 2011. CO emission per kWh assumes 1/3 of one percent of C emissions is CO based on Energy Information Administration 1994. PM10 emission per kWh from Layton 2004.
- CO2, NOx, SO2, and CO emission per Btu for natural gas, propane and butane (average used to represent LPG), Fuel #4 and #6 (average used to represent fuel oil and kerosene) from Leonardo Academy 2011.
- CO2 emissions per Btu of wood from Energy Information Administration 2014.
- CO, NOx and SOx emission per Btu based on total emissions and wood burning (tons) from (British Columbia Ministry 2005; Georgia Forestry Commission 2009).

Appendix II. Relative Tree Effects

The urban forest in Lake Forest Park Plot Inventory 2023 provides benefits that include carbon storage and sequestration, and air pollutant removal. To estimate the relative value of these benefits, tree benefits were compared to estimates of average municipal carbon emissions, average passenger automobile emissions, and average household emissions. See Appendix I for methodology.

Carbon storage is equivalent to:

- Annual carbon (C) emissions from 68,800 automobiles
- Annual C emissions from 28,200 single-family houses

Carbon monoxide removal is equivalent to:

- Annual carbon monoxide emissions from 7 automobiles
- Annual carbon monoxide emissions from 20 single-family houses

Nitrogen dioxide removal is equivalent to:

- Annual nitrogen dioxide emissions from 2,360 automobiles
- Annual nitrogen dioxide emissions from 1,060 single-family houses

Sulfur dioxide removal is equivalent to:

- Annual sulfur dioxide emissions from 14,500 automobiles
- Annual sulfur dioxide emissions from 38 single-family houses

Annual carbon sequestration is equivalent to:

- Annual C emissions from 1,900 automobiles
- Annual C emissions from 800 single-family houses

Appendix III. Comparison of Urban Forests

A common question asked is, "How does this city compare to other cities?" Although comparison among cities should be made with caution as there are many attributes of a city that affect urban forest structure and functions, summary data are provided from other cities analyzed using the i-Tree Eco model.

I. City totals for trees

City	% Tree Cover	Number of Trees	Carbon Storage (tons)	Carbon Sequestration (tons/yr)	Pollution Removal (tons/yr)
Toronto, ON, Canada	26.6	10,220,000	1,221,000	51,500	2,099
Atlanta, GA	36.7	9,415,000	1,344,000	46,400	1,663
Los Angeles, CA	11.1	5,993,000	1,269,000	77,000	1,975
New York, NY	20.9	5,212,000	1,350,000	42,300	1,676
London, ON, Canada	24.7	4,376,000	396,000	13,700	408
Chicago, IL	17.2	3,585,000	716,000	25,200	888
Phoenix, AZ	9.0	3,166,000	315,000	32,800	563
Baltimore, MD	21.0	2,479,000	570,000	18,400	430
Philadelphia, PA	15.7	2,113,000	530,000	16,100	575
Washington, DC	28.6	1,928,000	525,000	16,200	418
Oakville, ON, Canada	29.1	1,908,000	147,000	6,600	190
Albuquerque, NM	14.3	1,846,000	332,000	10,600	248
Boston, MA	22.3	1,183,000	319,000	10,500	283
Syracuse, NY	26.9	1,088,000	183,000	5,900	109
Woodbridge, NJ	29.5	986,000	160,000	5,600	210
Minneapolis, MN	26.4	979,000	250,000	8,900	305
San Francisco, CA	11.9	668,000	194,000	5,100	141
Morgantown, WV	35.5	658,000	93,000	2,900	72
Moorestown, NJ	28.0	583,000	117,000	3,800	118
Hartford, CT	25.9	568,000	143,000	4,300	58
Jersey City, NJ	11.5	136,000	21,000	890	41
Casper, WY	8.9	123,000	37,000	1,200	37
Freehold, NJ	34.4	48,000	20,000	540	22

II. Totals per acre of land area

City	Number of Trees/ac	Carbon Storage (tons/ac)	Carbon Sequestration (tons/ac/yr)	Pollution Removal (lb/ac/yr)
Toronto, ON, Canada	64.9	7.8	0.33	26.7
Atlanta, GA	111.6	15.9	0.55	39.4
Los Angeles, CA	19.6	4.2	0.16	13.1
New York, NY	26.4	6.8	0.21	17.0
London, ON, Canada	75.1	6.8	0.24	14.0
Chicago, IL	24.2	4.8	0.17	12.0
Phoenix, AZ	12.9	1.3	0.13	4.6
Baltimore, MD	48.0	11.1	0.36	16.6
Philadelphia, PA	25.1	6.3	0.19	13.6
Washington, DC	49.0	13.3	0.41	21.2
Oakville, ON, Canada	78.1	6.0	0.27	11.0
Albuquerque, NM	21.8	3.9	0.12	5.9
Boston, MA	33.5	9.1	0.30	16.1
Syracuse, NY	67.7	10.3	0.34	13.6
Woodbridge, NJ	66.5	10.8	0.38	28.4
Minneapolis, MN	26.2	6.7	0.24	16.3
San Francisco, CA	22.5	6.6	0.17	9.5
Morgantown, WV	119.2	16.8	0.52	26.0
Moorestown, NJ	62.1	12.4	0.40	25.1
Hartford, CT	50.4	12.7	0.38	10.2
Jersey City, NJ	14.4	2.2	0.09	8.6
Casper, WY	9.1	2.8	0.09	5.5
Freehold, NJ	38.3	16.0	0.44	35.3

Appendix IV. General Recommendations for Air Quality Improvement

Urban vegetation can directly and indirectly affect local and regional air quality by altering the urban atmosphere environment. Four main ways that urban trees affect air quality are (Nowak 1995):

- Temperature reduction and other microclimate effects
- Removal of air pollutants
- Emission of volatile organic compounds (VOC) and tree maintenance emissions
- Energy effects on buildings

The cumulative and interactive effects of trees on climate, pollution removal, and VOC and power plant emissions determine the impact of trees on air pollution. Cumulative studies involving urban tree impacts on ozone have revealed that increased urban canopy cover, particularly with low VOC emitting species, leads to reduced ozone concentrations in cities (Nowak 2000). Local urban management decisions also can help improve air quality.

Urban forest management strategies to help improve air quality include (Nowak 2000):

Strategy	Result
Increase the number of healthy trees	Increase pollution removal
Sustain existing tree cover	Maintain pollution removal levels
Maximize use of low VOC-emitting trees	Reduces ozone and carbon monoxide formation
Sustain large, healthy trees	Large trees have greatest per-tree effects
Use long-lived trees	Reduce long-term pollutant emissions from planting and removal
Use low maintenance trees	Reduce pollutants emissions from maintenance activities
Reduce fossil fuel use in maintaining vegetation	Reduce pollutant emissions
Plant trees in energy conserving locations	Reduce pollutant emissions from power plants
Plant trees to shade parked cars	Reduce vehicular VOC emissions
Supply ample water to vegetation	Enhance pollution removal and temperature reduction
Plant trees in polluted or heavily populated areas	Maximizes tree air quality benefits
Avoid pollutant-sensitive species	Improve tree health
Utilize evergreen trees for particulate matter	Year-round removal of particles

Appendix V. Tree Population

Species	Number of Trees	Percent of Population
Douglas fir (<i>Pseudotsuga menziesii</i>)	48,440	16.3%
Bigleaf maple (<i>Acer macrophyllum</i>)	32,880	11.1%
Western red cedar (<i>Thuja plicata</i>)	27,158	9.1%
Cherry laurel (<i>Prunus laurocerasus</i>)	24,377	8.2%
Bitter cherry (<i>Prunus emarginata</i>)	18,586	6.3%
English holly (<i>Ilex aquifolium</i>)	18,444	6.2%
Vine maple (<i>Acer circinatum</i>)	15,552	5.2%
European bird cherry (<i>Prunus padus</i>)	14,503	4.9%
Hinoki cypress (<i>Chamaecyparis obtusa</i>)	9,274	3.1%
Japanese maple (<i>Acer palmatum</i>)	8,406	2.8%
Western hemlock (<i>Tsuga heterophylla</i>)	8,368	2.8%
Northern white cedar (<i>Thuja occidentalis</i>)	6,744	2.3%
Red alder (<i>Alnus rubra</i>)	6,489	2.2%
Portugal laurel (<i>Prunus lusitanica</i>)	5,544	1.9%
Pacific dogwood (<i>Cornus nuttallii</i>)	4,637	1.6%
Plum spp (<i>Prunus</i>)	3,978	1.3%
Black poplar (<i>Populus nigra</i>)	3,626	1.2%
Swiss mountain pine (<i>Pinus mugo</i>)	2,719	0.9%
Camellia (<i>Camellia japonica</i>)	2,442	0.8%
Paper birch (<i>Betula papyrifera</i>)	2,023	0.7%
Lodgepole pine (<i>Pinus contorta</i>)	2,023	0.7%
Oneseed hawthorn (<i>Crataegus monogyna</i>)	1,918	0.6%
California laurel (<i>Umbellularia californica</i>)	1,813	0.6%
Cherry plum (<i>Prunus cerasifera</i>)	1,746	0.6%
Port orford cedar (<i>Chamaecyparis lawsoniana</i>)	1,641	0.6%
Kousa dogwood (<i>Cornus kousa</i>)	1,536	0.5%
Leyland cypress (<i>x Hesperotropis leylandii</i>)	1,469	0.5%
Western white pine (<i>Pinus monticola</i>)	1,221	0.4%
Blue spruce (<i>Picea pungens</i>)	1,116	0.4%
Callery pear (<i>Pyrus calleryana</i>)	1,116	0.4%
Deodar cedar (<i>Cedrus deodara</i>)	1,011	0.3%
Sitka spruce (<i>Picea sitchensis</i>)	1,011	0.3%
Giant Sequoia (<i>Sequoiadendron giganteum</i>)	1,011	0.3%
Flowering dogwood (<i>Cornus florida</i>)	954	0.3%
Red maple (<i>Acer rubrum</i>)	906	0.3%
Japanese angelica tree (<i>Aralia elata</i>)	906	0.3%
Atlas cedar (<i>Cedrus atlantica</i>)	906	0.3%
Katsura tree (<i>Cercidiphyllum japonicum</i>)	906	0.3%
Blue chinese fir (<i>Cunninghamia lanceolata</i>)	906	0.3%
Chinese parasoltree (<i>Firmiana simplex</i>)	906	0.3%

Cucumber tree (<i>Magnolia acuminata</i>)	906	0.3%
Babylon weeping willow (<i>Salix babylonica</i>)	906	0.3%
Pacific yew (<i>Taxus brevifolia</i>)	906	0.3%
Apple spp (<i>Malus</i>)	839	0.3%
Sweet cherry (<i>Prunus avium</i>)	629	0.2%
European white birch (<i>Betula pendula</i>)	315	0.1%
Southern magnolia (<i>Magnolia grandiflora</i>)	315	0.1%
Japanese flowering cherry (<i>Prunus serrulata</i>)	315	0.1%
Common apple (<i>Malus domestica</i>)	210	<0.1%
Black locust (<i>Robinia pseudoacacia</i>)	210	<0.1%
European mountain ash (<i>Sorbus aucuparia</i>)	210	<0.1%
Subalpine fir (<i>Abies lasiocarpa</i>)	105	<0.1%
Trident maple (<i>Acer buergerianum</i>)	105	<0.1%
Eastern redbud (<i>Cercis canadensis</i>)	105	<0.1%
Oregon ash (<i>Fraxinus latifolia</i>)	105	<0.1%
Honeylocust (<i>Gleditsia triacanthos</i>)	105	<0.1%
Black walnut (<i>Juglans nigra</i>)	105	<0.1%
Golden Chain Tree spp (<i>Laburnum</i>)	105	<0.1%
Lagerstroemia spp (<i>Lagerstroemia</i>)	105	<0.1%
Privet spp (<i>Ligustrum</i>)	105	<0.1%
Norway spruce (<i>Picea abies</i>)	105	<0.1%
Scots pine (<i>Pinus sylvestris</i>)	105	<0.1%
Black cottonwood (<i>Populus balsamifera</i> ssp.	105	<0.1%
Common plum (<i>Prunus domestica</i>)	105	<0.1%
Sargent cherry (<i>Prunus sargentii</i>)	105	<0.1%
Umbrella pine (<i>Sciadopitys verticillata</i>)	105	<0.1%
Coast redwood (<i>Sequoia sempervirens</i>)	105	<0.1%
Yew spp (<i>Taxus</i>)	105	<0.1%
Mountain hemlock (<i>Tsuga mertensiana</i>)	105	<0.1%
Sweetgum (<i>Liquidambar styraciflua</i>)	96	<0.1%
River birch (<i>Betula nigra</i>)	72	<0.1%
Willow spp (<i>Salix</i>)	24	<0.1%
Total	297,056	100%





Appendix VI. Potential Risk of Pests

Fifty-three insects and diseases were analyzed to quantify their potential impact on the urban forest. As each insect/disease is likely to attack different host tree species, the implications for {0} will vary. The number of trees at risk reflects only the known host species that are likely to experience mortality.

Code	Scientific Name	Common Name	Trees at Risk (#)	Value (\$ millions)
AL	<i>Phyllocnistis populiella</i>	Aspen Leafminer	4,107	2.35
ALB	<i>Anoplophora glabripennis</i>	Asian Longhorned Beetle	66,215	125.94
ARCA	<i>Neodothiora populina</i>	Aspen Running Canker	0	0.00
ARD	<i>Armillaria</i> spp.	Armillaria Root Disease	75,703	266.33
BBD	<i>Neonectria faginata</i>	Beech Bark Disease	0	0.00
BC	<i>Sirococcus clavigignenti</i> <i>juglandacearum</i>	Butternut Canker	105	0.02
BLD	<i>Litylenchus crenatae</i> mccannii	Beech Leaf Disease	0	0.00
BM	<i>Euproctis chrysorrhoea</i>	Browntail Moth	6,864	3.78
BOB	<i>Tubakia iowensis</i>	Bur Oak Blight	0	0.00
BSRD	<i>Leptographium wageneri</i>	Black Stain Root Disease	60,157	203.03
BWA	<i>Adelges piceae</i>	Balsam Woolly Adelgid	210	1.26
CB	<i>Cryphonectria parasitica</i>	Chestnut Blight	0	0.00
DA	<i>Discula destructiva</i>	Dogwood Anthracnose	7,127	2.27
DBSR	<i>Leptographium wageneri</i> var. <i>pseudotsugae</i>	Douglas-fir Black Stain Root Disease	60,157	203.03
DED	<i>Ophiostoma novo-ulmi</i>	Dutch Elm Disease	0	0.00
DFB	<i>Dendroctonus pseudotsugae</i>	Douglas-Fir Beetle	48,440	185.81
EAB	<i>Agrilus planipennis</i>	Emerald Ash Borer	105	0.23
FE	<i>Scolytus ventralis</i>	Fir Engraver	48,650	185.96
FR	<i>Cronartium quercuum</i> f. sp. Fusiforme	Fusiform Rust	0	0.00
FTC	<i>Malacosoma disstria</i>	Forest Tent Caterpillar	10,491	21.72
GM	<i>Lymantria dispar</i>	Gypsy Moth	14,913	35.40
GSOB	<i>Agrilus auroguttatus</i>	Goldspotted Oak Borer	0	0.00
HRD	<i>Heterobasidion irregulare/</i> <i>occidentale</i>	Heterobasidion Root Disease	87,420	284.66
HS	<i>Neodiprion tsugae</i>	Hemlock Sawfly	9,484	12.18
HWA	<i>Adelges tsugae</i>	Hemlock Woolly Adelgid	0	0.00
JPB	<i>Dendroctonus jeffreyi</i>	Jeffrey Pine Beetle	0	0.00
JPBW	<i>Choristoneura pinus</i>	Jack Pine Budworm	4,847	1.44
LAT	<i>Choristoneura conflictana</i>	Large Aspen Tortrix	13,005	24.15
LWD	<i>Raffaelea lauricola</i>	Laurel Wilt	1,813	0.55
MOB	<i>Xyleborus monographus</i>	Mediterranean Oak Borer	629	0.38
MPB	<i>Dendroctonus ponderosae</i>	Mountain Pine Beetle	3,454	8.95
NSE	<i>Ips perturbatus</i>	Northern Spruce Engraver	1,011	2.68
OW	<i>Ceratocystis fagacearum</i>	Oak Wilt	0	0.00
PBSR	<i>Leptographium wageneri</i> var. <i>ponderosum</i>	Pine Black Stain Root Disease	2,023	1.30
POCRD	<i>Phytophthora lateralis</i>	Port-Orford-Cedar Root Disease	18,565	8.64

Code	Scientific Name	Common Name	Trees at Risk (#)	Value (\$ millions)
PSB	<i>Tomicus piniperda</i>	Pine Shoot Beetle	54,613	194.90
PSHB	<i>Euwallacea nov. sp.</i>	Polyphagous Shot Hole Borer	54,461	134.92
RPS	<i>Matsucoccus resinosae</i>	Red Pine Scale	0	0.00
SB	<i>Dendroctonus rufipennis</i>	Spruce Beetle	2,233	5.58
SBW	<i>Choristoneura fumiferana</i>	Spruce Budworm	49,661	188.71
SFM	subalpine fir mortality summary	Subalpine Fir Mortality	105	0.05
SLF	<i>Lycorma delicatula</i>	Spotted Lanternfly	22,400	12.08
SOD	<i>Phytophthora ramorum</i>	Sudden Oak Death	87,875	298.44
SPB	<i>Dendroctonus frontalis</i>	Southern Pine Beetle	16,773	22.95
SW	<i>Sirex noctilio</i>	Sirex Wood Wasp	6,173	9.09
TCD	<i>Geosmithia morbida</i>	Thousand Canker Disease	105	0.02
WBB	<i>Dryocoetes confusus</i>	Western Bark Beetle	0	0.00
WBBU	<i>Acleris gloverana</i>	Western Blackheaded Budworm	56,913	195.30
WFNPM	western five-needle pine mortality summary	Western Five-Needle Pine Mortality	1,221	6.43
WM	<i>Operophtera brumata</i>	Winter Moth	61,966	132.94
WPB	<i>Dendroctonus brevicomis</i>	Western Pine Beetle	0	0.00
WPBR	<i>Cronartium ribicola</i>	White Pine Blister Rust	1,221	6.43
WSB	<i>Choristoneura occidentalis</i>	Western Spruce Budworm	59,565	204.70

In the following graph, the pests are color coded according to the county's proximity to the pest occurrence in the United States. Red indicates that the pest is within the county; orange indicates that the pest is within 250 miles of the county; yellow indicates that the pest is within 750 miles of the county; and green indicates that the pest is outside of these ranges.



Note: points - Number of trees, bars - Replacement value

Based on the host tree species for each pest and the current range of the pest (Forest Health Technology Enterprise Team 2014), it is possible to determine what the risk is that each tree species in the urban forest could be attacked by an insect or disease.

Spp. Risk	Species Name	AL	ALB	ARCA	ARD	BBD	BC	BLD	BM	BOB	BSRD	BWA	CB	DA	DBSR	DED	DFB	EAB	FE	FR	FTC	GM	GSOB	HRD	HS	HWA	JPB	JPBW	LAT	LWD	MOB	MPB	NSE	OW
32	Douglas-fir																																	
29	Western white pine																																	
24	Subalpine fir																																	
24	Norway spruce																																	
21	Lodgepole pine																																	
19	Western hemlock																																	
19	Mountain hemlock																																	
16	Willow spp																																	
12	Plum spp																																	
12	Scots pine																																	
12	Black cottonwood																																	
11	Paper birch																																	
11	Sitka spruce																																	
10	Red alder																																	
10	Blue spruce																																	
10	European white birch																																	
10	River birch																																	
8	Western red cedar																																	
6	Apple spp																																	
5	Bigleaf maple																																	
5	Arborvitae																																	
5	Black poplar																																	
5	Black walnut																																	
4	Hinoki cypress																																	
4	Swiss mountain pine																																	
4	California laurel																																	
4	Port orford cedar																																	
4	Callery pear																																	
4	Pacific yew																																	
4	Sweet cherry																																	
4	European mountain ash																																	
4	Oregon ash																																	
4	Sweetgum																																	
3	Japanese maple																																	
3	Pacific dogwood																																	
3	Red maple																																	
3	Trident maple																																	
2	Chinese parasoltree																																	
2	Japanese flowering cherry																																	

Spp. Risk	Risk Weight	Species Name	AL	ALB	ARCA	ARD	BBD	BC	BLD	BM	BOB	BSRD	BWA	CB	DA	DBSR	DED	DFB	EAB	FE	FR	FTC	GM	GSOB	HRD	HS	HWA	JPB	JPBW	LAT	LWD	MOB	MPB	NSE	OW
2		Coast redwood																																	
1		Bitter cherry																																	
1		Vine maple																																	
1		Camellia																																	
1		Common hawthorn																																	
1		Kousa dogwood																																	
1		Flowering dogwood																																	
1		Japanese angelica tree																																	
1		Atlas cedar																																	
1		Katsura tree																																	
1		Babylon weeping willow																																	
1		Southern magnolia																																	
1		Black locust																																	
1		Honeylocust																																	
1		Common plum																																	

Spp. Risk	Risk Weight	Species Name	PBSR	POCRD	PSB	PSHB	RPS	SB	SBW	SFM	SLF	SOD	SPB	SW	TCD	WBB	WBBU	WFNPW	WM	WPB	WPBR	WSB
32		Douglas-fir																				
29		Western white pine																				
24		Subalpine fir																				
24		Norway spruce																				
21		Lodgepole pine																				
19		Western hemlock																				
19		Mountain hemlock																				
16		Willow spp																				
12		Plum spp																				
12		Scots pine																				
12		Black cottonwood																				
11		Paper birch																				
11		Sitka spruce																				
10		Red alder																				
10		Blue spruce																				
10		European white birch																				
10		River birch																				
8		Western red cedar																				
6		Apple spp																				
5		Bigleaf maple																				
5		Arborvitae																				
5		Black poplar																				
5		Black walnut																				
4		Hinoki cypress																				
4		Swiss mountain pine																				
4		California laurel																				
4		Port orford cedar																				

Spp. Risk	Risk Weight	Species Name	PBSR	POCRD	PSB	PSHB	RPS	SB	SBW	SFM	SLF	SOD	SPB	SW	TCD	WBB	WBBU	WFNPM	WM	WPB	WPBR	WSB
4	4	Callery pear																				
4	4	Pacific yew					Red															
4	4	Sweet cherry											Green									
4	4	European mountain ash																				
4	4	Oregon ash																		Green		
4	4	Sweetgum																				
3	3	Japanese maple																				
3	3	Pacific dogwood																Yellow				
3	3	Red maple																				
3	3	Trident maple																				
2	2	Chinese parasoltree																				
2	2	Japanese flowering cherry																				
2	2	Coast redwood																Yellow				
1	1	Bitter cherry																				
1	1	Vine maple																		Green		
1	1	Camellia																				
1	1	Common hawthorn																	Green			
1	1	Kousa dogwood																				
1	1	Flowering dogwood																				
1	1	Japanese angelica tree																Green				
1	1	Atlas cedar																				
1	1	Katsura tree																				
1	1	Babylon weeping willow																				
1	1	Southern magnolia																				
1	1	Black locust																Green				
1	1	Honeylocust																				
1	1	Common plum																	Green			

Note:

Species that are not listed in the matrix are not known to be hosts to any of the pests analyzed.

Species Risk:

- Red indicates that tree species is at risk to at least one pest within county
- Orange indicates that tree species has no risk to pests in county, but has a risk to at least one pest within 250 miles from the county
- Yellow indicates that tree species has no risk to pests within 250 miles of county, but has a risk to at least one pest that is 250 and 750 miles from the county
- Green indicates that tree species has no risk to pests within 750 miles of county, but has a risk to at least one pest that is greater than 750 miles from the county

Risk Weight:

Numerical scoring system based on sum of points assigned to pest risks for species. Each pest that could attack tree species is scored as 4 points if red, 3 points if orange, 2 points if yellow and 1 point if green.

Pest Color Codes:

- Red indicates pest is within King county
- Red indicates pest is within 250 miles county
- Yellow indicates pest is within 750 miles of King county
- Green indicates pest is outside of these ranges

References for i-Tree Report and Appendices

Abdollahi, K.K.; Ning, Z.H.; Appeaning, A., eds. 2000. Global climate change and the urban forest. Baton Rouge, LA: GCRCC and Franklin Press. 77 p.

Baldocchi, D. 1988. A multi-layer model for estimating sulfur dioxide deposition to a deciduous oak forest canopy. *Atmospheric Environment*. 22: 869-884.

Baldocchi, D.D.; Hicks, B.B.; Camara, P. 1987. A canopy stomatal resistance model for gaseous deposition to vegetated surfaces. *Atmospheric Environment*. 21: 91-101.

Bidwell, R.G.S.; Fraser, D.E. 1972. Carbon monoxide uptake and metabolism by leaves. *Canadian Journal of Botany*. 50: 1435-1439.

British Columbia Ministry of Water, Land, and Air Protection. 2005. Residential wood burning emissions in British Columbia. British Columbia.

Broecker, W.S. 1970. Man's oxygen reserve. *Science* 168(3939): 1537-1538.

Bureau of Transportation Statistics. 2010. Estimated National Average Vehicle Emissions Rates per Vehicle by Vehicle Type using Gasoline and Diesel. Washington, DC: Bureau of Transportation Statistics, U.S. Department of Transportation. Table 4-43.

California Air Resources Board. 2013. Methods to Find the Cost-Effectiveness of Funding Air Quality Projects. Table 3 Average Auto Emission Factors. CA: California Environmental Protection Agency, Air Resources Board.

Carbon Dioxide Information Analysis Center. 2010. CO₂ Emissions (metric tons per capita). Washington, DC: The World Bank.

Cardelino, C.A.; Chameides, W.L. 1990. Natural hydrocarbons, urbanization, and urban ozone. *Journal of Geophysical Research*. 95(D9): 13,971-13,979.

Ciesla, W. M.; Kruse, J. J. 2009. Large Aspen Tortrix. Forest Insect & Disease Leaflet 139. Washington, DC: U. S. Department of Agriculture, Forest Service. 8 p.

DeMars, C. J., Jr.; Roettgering, B. H. 1982. Western Pine Beetle. Forest Insect & Disease Leaflet 1. Washington, DC: U.S. Department of Agriculture, Forest Service. 8 p.

Eastern Forest Environmental Threat Assessment Center. Dutch Elm Disease. <http://threatsummary.forestthreats.org/>

Energy Information Administration. 1994. Energy Use and Carbon Emissions: Non-OECD Countries. Washington, DC: Energy Information Administration, U.S. Department of Energy.

Energy Information Administration. 2013. CE2.1 Fuel consumption totals and averages, U.S. homes. Washington, DC: Energy Information Administration, U.S. Department of Energy.

Energy Information Administration. 2014. CE5.2 Household wood consumption. Washington, DC: Energy Information Administration, U.S. Department of Energy.

Federal Highway Administration. 2013. Highway Statistics 2011. Washington, DC: Federal Highway Administration, U.S. Department of Transportation. Table VM-1.

Fellin, D. G.; Dewey, J. E. 1986. Western Spruce Budworm. Forest Insect & Disease Leaflet 53. Washington, DC: U.S. Department of Agriculture, Forest Service. 10 p.

Ferrell, G. T. 1986. Fir Engraver. Forest Insect & Disease Leaflet 13. Washington, DC: U. S. Department of Agriculture, Forest Service. 8 p.

Forest Health Technology Enterprise Team. 2014. 2012 National Insect & Disease Risk Maps/Data. Fort Collins, CO: U.S. Department of Agriculture, Forest Service. <http://www.fs.fed.us/foresthealth/technology/nidrm2012.shtml>

Georgia Forestry Commission. 2009. Biomass Energy Conversion for Electricity and Pellets Worksheet. Dry Branch, GA: Georgia Forestry Commission.

Gibson, K.; Kegley, S.; Bentz, B. 2009. Mountain Pine Beetle. Forest Insect & Disease Leaflet 2. Washington, DC: U. S. Department of Agriculture, Forest Service. 12 p.

Heirigs, P.L.; Delaney, S.S.; Dulla, R.G. 2004. Evaluation of MOBILE Models: MOBILE6.1 (PM), MOBILE6.2 (Toxics), and MOBILE6/CNG. Sacramento, CA: National Cooperative Highway Research Program, Transportation Research Board.

Hirabayashi, S. 2011. Urban Forest Effects-Dry Deposition (UFORE-D) Model Enhancements, <http://www.itreetools.org/eco/resources/UFORE-D%20enhancements.pdf>

Hirabayashi, S. 2012. i-Tree Eco Precipitation Interception Model Descriptions, http://www.itreetools.org/eco/resources/iTree_Eco_Precipitation_Interception_Model_Descriptions_V1_2.pdf

Hirabayashi, S.; Kroll, C.; Nowak, D. 2011. Component-based development and sensitivity analyses of an air pollutant dry deposition model. Environmental Modeling and Software. 26(6): 804-816.

Hirabayashi, S.; Kroll, C.; Nowak, D. 2012. i-Tree Eco Dry Deposition Model Descriptions V 1.0

Holsten, E.H.; Thier, R.W.; Munson, A.S.; Gibson, K.E. 1999. The Spruce Beetle. Forest Insect & Disease Leaflet 127. Washington, DC: U.S. Department of Agriculture, Forest Service. 12 p.

Interagency Working Group on Social Cost of Carbon, United States Government. 2015. Technical Support Document: Technical Update of the Social Cost of Carbon for Regulatory Impact Analysis Under Executive Order 12866. <http://www.whitehouse.gov/sites/default/files/omb/info/reg/scc-tsd-final-july-2015.pdf>

Layton, M. 2004. 2005 Electricity Environmental Performance Report: Electricity Generation and Air Emissions. CA:

California Energy Commission.

Leonardo Academy. 2011. Leonardo Academy's Guide to Calculating Emissions Including Emission Factors and Energy Prices. Madison, WI: Leonardo Academy Inc.

Liebhold, A. 2010 draft. Personal communication on the geographic distribution of forest pest species.

Lovett, G.M. 1994. Atmospheric deposition of nutrients and pollutants in North America: an ecological perspective. *Ecological Applications*. 4: 629-650.

McPherson, E.G.; Maco, S.E.; Simpson, J.R.; Peper, P.J.; Xiao, Q.; VanDerZanden, A.M.; Bell, N. 2002. Western Washington and Oregon Community Tree Guide: Benefits, Costs, and Strategic Planting. International Society of Arboriculture, Pacific Northwest, Silverton, OR.

McPherson, E.G.; Simpson, J.R. 1999. Carbon dioxide reduction through urban forestry: guidelines for professional and volunteer tree planters. Gen. Tech. Rep. PSW-171. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. 237 p.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Crowell, A.M.N.; Xiao, Q. 2010. Northern California coast community tree guide: benefits, costs, and strategic planting. PSW-GTR-228. Gen. Tech. Rep. PSW-GTR-228. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Gardner, S.L.; Vargas, K.E.; Maco, S.E.; Xiao, Q. 2006a. Coastal Plain Community Tree Guide: Benefits, Costs, and Strategic Planting PSW-GTR-201. USDA Forest Service, Pacific Southwest Research Station, Albany, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Gardner, S.L.; Vargas, K.E.; Xiao, Q. 2007. Northeast community tree guide: benefits, costs, and strategic planting.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Maco, S.E.; Gardner, S.L.; Cozad, S.K.; Xiao, Q. 2006b. Midwest Community Tree Guide: Benefits, Costs and Strategic Planting PSW-GTR-199. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Maco, S.E.; Gardner, S.L.; Vargas, K.E.; Xiao, Q. 2006c. Piedmont Community Tree Guide: Benefits, Costs, and Strategic Planting PSW-GTR 200. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Maco, S.E.; Xiao Q.; Mulrean, E. 2004. Desert Southwest Community Tree Guide: Benefits, Costs and Strategic Planting. Phoenix, AZ: Arizona Community Tree Council, Inc. 81 :81.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Scott, K.I.; Xiao, Q. 2000. Tree Guidelines for Coastal Southern California Communities. Local Government Commission, Sacramento, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Xiao, Q. 1999. Tree Guidelines for San Joaquin Valley Communities. Local Government Commission, Sacramento, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Xiao, Q.; Maco, S.E.; Hoefer, P.J. 2003. Northern Mountain and Prairie Community Tree Guide: Benefits, Costs and Strategic Planting. Center for Urban Forest Research, USDA Forest Service, Pacific Southwest Research Station, Albany, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Xiao, Q.; Pittenger, D.R.; Hodel, D.R. 2001. Tree Guidelines for Inland

Empire Communities. Local Government Commission, Sacramento, CA.

Murray, F.J.; Marsh L.; Bradford, P.A. 1994. New York State Energy Plan, vol. II: issue reports. Albany, NY: New York State Energy Office.

National Invasive Species Information Center. 2011. Beltsville, MD: U.S. Department of Agriculture, National Invasive Species Information Center. <http://www.invasivespeciesinfo.gov/plants/main.shtml>

Nicholls, T. H.; Anderson, R. L. 1977. How to Identify White Pine Blister Rust and Remove Cankers. St. Paul, MN: U.S. Department of Agriculture, Forest Service, Northeastern Area State and Private Forestry

Nowak, D.J. 1994. Atmospheric carbon dioxide reduction by Chicago's urban forest. In: McPherson, E.G.; Nowak, D.J.; Rountree, R.A., eds. Chicago's urban forest ecosystem: results of the Chicago Urban Forest Climate Project. Gen. Tech. Rep. NE-186. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station: 83-94.

Nowak, D.J. 1995. Trees pollute? A "TREE" explains it all. In: Proceedings of the 7th National Urban Forestry Conference. Washington, DC: American Forests: 28-30.

Nowak, D.J. 2000. The interactions between urban forests and global climate change. In: Abdollahi, K.K.; Ning, Z.H.; Appeaning, A., eds. Global Climate Change and the Urban Forest. Baton Rouge, LA: GCRCC and Franklin Press: 31-44.

Nowak, D.J., Hirabayashi, S., Bodine, A., Greenfield, E. 2014. Tree and forest effects on air quality and human health in the United States. Environmental Pollution. 193:119-129.

Nowak, D.J., Hirabayashi, S., Bodine, A., Hoehn, R. 2013. Modeled PM2.5 removal by trees in ten U.S. cities and associated health effects. Environmental Pollution. 178: 395-402.

Nowak, D.J.; Civerolo, K.L.; Rao, S.T.; Sistla, S.; Luley, C.J.; Crane, D.E. 2000. A modeling study of the impact of urban trees on ozone. Atmospheric Environment. 34: 1601-1613.

Nowak, D.J.; Crane, D.E. 2000. The Urban Forest Effects (UFORE) Model: quantifying urban forest structure and functions. In: Hansen, M.; Burk, T., eds. Integrated tools for natural resources inventories in the 21st century. Proceedings of IUFRO conference. Gen. Tech. Rep. NC-212. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Research Station: 714-720.

Nowak, D.J.; Crane, D.E.; Dwyer, J.F. 2002a. Compensatory value of urban trees in the United States. Journal of Arboriculture. 28(4): 194 - 199.

Nowak, D.J.; Crane, D.E.; Stevens, J.C.; Hoehn, R.E. 2005. The urban forest effects (UFORE) model: field data collection manual. V1b. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Research Station, 34 p. http://www.fs.fed.us/ne/syracuse/Tools/downloads/UFORE_Manual.pdf

Nowak, D.J.; Crane, D.E.; Stevens, J.C.; Ibarra, M. 2002b. Brooklyn's urban forest. Gen. Tech. Rep. NE-290. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Research Station. 107 p.

Nowak, D.J.; Dwyer, J.F. 2000. Understanding the benefits and costs of urban forest ecosystems. In: Kuser, John, ed. Handbook of urban and community forestry in the northeast. New York, NY: Kluwer Academic/Plenum: 11-22.

Nowak, D.J.; Hoehn, R.; Crane, D. 2007. Oxygen production by urban trees in the United States. Arboriculture & Urban Forestry. 33(3):220-226.

Nowak, D.J.; Hoehn, R.E.; Crane, D.E.; Stevens, J.C.; Walton, J.T; Bond, J. 2008. A ground-based method of assessing urban forest structure and ecosystem services. *Arboriculture and Urban Forestry*. 34(6): 347-358.

Nowak, D.J.; Stevens, J.C.; Sisinni, S.M.; Luley, C.J. 2002c. Effects of urban tree management and species selection on atmospheric carbon dioxide. *Journal of Arboriculture*. 28(3): 113-122.

Oregon Invasive Species Council. 2014. 100 Worst List. OR: Oregon Invasive Species Council. <http://www.oregoninvasivespeciescouncil.org/media/100_worst.pdf>

Peper, P.J.; McPherson, E.G.; Simpson, J.R.; Albers, S.N.; Xiao, Q. 2010. Central Florida community tree guide: benefits, costs, and strategic planting. Gen. Tech. Rep. PSW-GTR-230. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

Peper, P.J.; McPherson, E.G.; Simpson, J.R.; Vargas, K.E.; Xiao Q. 2009. Lower Midwest community tree guide: benefits, costs, and strategic planting. PSW-GTR-219. Gen. Tech. Rep. PSW-GTR-219. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

Schmitz, R. F.; Gibson, K. E. 1996. Douglas-fir Beetle. Forest Insect & Disease Leaflet 5. R1-96-87. Washington, DC: U. S. Department of Agriculture, Forest Service. 8 p.

U.S. Environmental Protection Agency. 2010. Light-Duty Vehicle Greenhouse Gas Emission Standards and Corporate Average Fuel Economy Standards. Washington, DC: U.S. Environmental Protection Agency. EPA-420-R-10-012a

U.S. Environmental Protection Agency. 2015. The social cost of carbon. <http://www.epa.gov/climatechange/EPAactivities/economics/scc.html>

van Essen, H.; Schrotten, A.; Otten, M.; Sutter, D.; Schreyer, C.; Zandonella, R.; Maibach, M.; Doll, C. 2011. External Costs of Transport in Europe. Netherlands: CE Delft. 161 p.

Vargas, K.E.; McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Gardner, S.L.; Xiao, Q. 2007a. Interior West Tree Guide.

Vargas, K.E.; McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Gardner, S.L.; Xiao, Q. 2007b. Temperate Interior West Community Tree Guide: Benefits, Costs, and Strategic Planting.

Vargas, K.E.; McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Gardner, S.L.; Xiao, Q. 2008. Tropical community tree guide: benefits, costs, and strategic planting. PSW-GTR-216. Gen. Tech. Rep. PSW-GTR-216. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

Worrall, J.J. 2007. Chestnut Blight. Forest and Shade Tree Pathology. http://www.forestpathology.org/dis_chestnut.html

Zinke, P.J. 1967. Forest interception studies in the United States. In: Sopper, W.E.; Lull, H.W., eds. *Forest Hydrology*. Oxford, UK: Pergamon Press: 137-161.